

# Resonances and formation of the gaps in the spectrum of quasi-periodic Schrödinger equation

Michael Goldstein , University of Toronto

February 7, 2006

## Abstract

This talk is a review of some results on spectrum and localized eigen functions of quasi-periodic Schrödinger equation , obtained in recent joint works with Wilhelm Schlag. We consider one-dimensional difference Schrödinger equation

$$[H(x, \omega)\varphi](n) \equiv -\varphi(n-1) - \varphi(n+1) + V(x+n\omega)\varphi(n) = E\varphi(n) ,$$

$n \in \mathbb{Z}$ ,  $x, \omega \in [0, 1]$  with real analytic function  $V(x)$ . It was established in [GolSch2] that the positivity of the Lyapunov exponent for  $E$  belonging to some interval  $(E', E'')$  implies the absolute continuity of the integrated density of states on this interval for a.a. $\omega$ . In this talk I will explain how we apply the technology and the results of [GolSch2], to describe the mechanism responsible for formation of a dense in  $(E', E'')$  set of intervals free from the points of  $\bigcup_x \text{sp } H(x, \omega)$  in the regime of positive Lyapunov exponent. The results on gaps are effective in finite volume approximation , and therefore are both constructive and quantitative.

## References

- [Bou] Bourgain, J. *Hölder regularity of integrated density of states for the almost Mathieu operator in a perturbative regime*. Lett. Math. Phys. 51 (2000), no. 2, 83–118.
- [BouGol] Bourgain, J., Goldstein, M. *On nonperturbative localization with quasi-periodic potential*. Ann. of Math. (2) 152 (2000), no. 3, 835–879.
- [BouGolSch] Bourgain, J., Goldstein, M., Schlag, W. *Anderson localization for Schrödinger operators on  $\mathbb{Z}$  with potentials given by the skew-shift*. Comm. Math. Phys. 220 (2001), no. 3, 583–621.
- [FroSpe1] Fröhlich, J., Spencer, T. *Absence of diffusion in the Anderson tight binding model for large disorder or low energy*. Comm. Math. Phys. **88** (1983), 151–189.
- [FroSpeWit] Fröhlich, J., Spencer, T., Wittwer, P. *Localization for a class of one dimensional quasi-periodic Schrödinger operators*. Commun. Math. Phys. **132** (1990), 5–25.
- [GolSch1] Goldstein, M., Schlag, W. *Hölder continuity of the integrated density of states for quasiperiodic Schrödinger equations and averages of shifts of subharmonic functions*. Ann. of Math. (2) 154 (2001), no. 1, 155–203.

- [GolSch2] Goldstein, M., Schlag, W. *Five properties of the integrated density of states and quantitative separation property of Dirichlet eigenvalues*, Preprint.
- [Sin1] Sinai, Y. G. *Anderson localization for one-dimensional difference Schrödinger operator with quasi-periodic potential*. J. Stat. Phys. **46** (1987), 861–909.
- [Pui] Puig, J. *Cantor spectrum for the almost Mathieu operator*. Comm. Math. Phys. 224 (2004) , no. 2, 297-309.